


Ankylosing Spondylitis (AS)

AS is a chronic inflammatory disease, predominantly involving the spine and sacroiliac (SI) joints.¹⁻⁴ Patients with AS have clearly defined structural damage of SI joints on X-ray.^{3,4}



Onset and diagnosis

Affects young people: mean age of symptom onset = in mid-20s.⁵ Average delay of

8.5 years to diagnosis⁶

Prevalence


Global prevalence (%) of AS⁷



Estimated prevalence of AS in adults aged

18 to 44 years is 0.35%*⁸

Patients may experience additional disease-related manifestations^{9,10}



Symptoms^{2,11-13}



References:

1. van der Heijde D, et al. *Ann Rheum Dis.* 2017. pii: annrheumdis-2016-210770
 2. Rudwaleit M, et al. *Arthritis Rheum.* 2005;52:1000-8
 3. Sieper J, van der Heijde D. *Arth Rheum.* 2013;65:543-51
 4. Poddubnyy D, et al. *Ann Rheum Dis.* 2011;70:1369-74
 5. Sieper J, Rudwaleit M. *Ann Rheum Dis.* 2005;64:659-63
 6. Sykes MP, et al. *Rheumatology* 2015;54:2283-4
 7. Stolwijk C, et al. *Arthritis Care Res.* 2016;68:1320-31
 8. Strand V, et al. *Arthritis Care Res.* 2013;65:1299-306
 9. de Winter JJ, et al. *Arthritis Res Ther.* 2016;18:196
 10. Davey-Ranasinghe N, Deodhar A. *Curr Opin Rheumatol.* 2013;25:509-16
 11. Rudwaleit M, et al. *Arthritis Rheum.* 2006;54:569-78
 12. Dougados M, et al. *Clin Exp Rheumatol.* 2017;35:209-13
 13. Levermore S, et al. *Rheumatol Int.* 2017;37:257-71
- *Based on a study conducted in the United States.



HQ/0417/CI/00039a; Date of preparation: June 2017



Inspired by patients.
Driven by science.