

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/346018595>

Diagnostic delay in axial spondyloarthritis: a systematic review and meta-analysis

Article in *Rheumatology* · November 2020

DOI: 10.1093/rheumatology/keaa807

CITATIONS

0

READS

123

6 authors, including:



Sizheng Steven Zhao
University of Liverpool

116 PUBLICATIONS 503 CITATIONS

[SEE PROFILE](#)



Bradley Pittam
University Hospital Of South Manchester NHS Foundation Trust

9 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)



Ashar Ahmed
Southport and Ormskirk Hospital NHS Trust

11 PUBLICATIONS 13 CITATIONS

[SEE PROFILE](#)



Nicola J Goodson
University of Liverpool

162 PUBLICATIONS 3,270 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Birth month and adult bone mineral density [View project](#)



Effectiveness comparison of osteoporosis medications [View project](#)

Diagnostic delay in axial spondyloarthritis: a systematic review and meta-analysis

Sizheng Steven Zhao^{1,2}, Bradley Pittam³, Nicholas L Harrison², Ashar E Ahmed⁴, Nicola J Goodson², David M Hughes⁵

1 Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool

2 Department of Rheumatology, Liverpool University Hospitals, Liverpool

3 School of Medicine, University of Liverpool, Liverpool

4 Department of Rheumatology, Southport & Ormskirk Hospital, Southport

5 Department of Biostatistics, Institute of Translational Medicine, University of Liverpool, Liverpool, UK

Correspondence to:

Dr Sizheng S Zhao

Department of Rheumatology

Liverpool University Hospitals

Liverpool

L9 7AL

UK

s.zhao8@liverpool.ac.uk

Abstract

Background. Delay to diagnosis in axial spondyloarthritis (axSpA) is longer than many other rheumatic diseases. Prolonged delay is associated with poorer outcomes including functional impairment and quality of life. Our aims were to describe 1) global variation in delay to diagnosis, 2) factors associated with delay, and 3) delay compared with psoriatic arthritis (PsA).

Methods. We searched Medline, PubMed, EMBASE and Web of Science using a predefined protocol. Diagnostic delay was defined as years between ages at symptom onset and diagnosis. We pooled mean delay using random-effects inverse variance meta-analysis. We examined variations in pooled estimates using pre-specified subgroup analyses and sources of heterogeneity using meta-regression.

Results. A total of 64 studies reported mean diagnostic delay in axSpA patients. The pooled mean delay was 6.7 years (95%CI 6.2 to 7.2) with high levels of heterogeneity. Delay to diagnosis did not improve over time when stratifying results by year of publication. Studies from high-income countries (defined by the World Bank) reported longer delay than those from middle-income countries. Factors consistently reported to be associated with longer delay were: lower education levels, younger age at symptom onset and absence of extra-articular manifestations (EAMs). Pooled estimate for diagnostic delay from 8 PsA studies was significantly shorter, at 2.6 years (95%CI 1.6 to 3.6).

Conclusion. For axSpA patients, delay to diagnosis remains unacceptably prolonged in many parts of the world. Patient factors (e.g., education) and disease presentation (onset age and EAMs) should inform campaigns to improve delay.

Keywords: axial spondyloarthritis, ankylosing spondylitis, delay to diagnosis, psoriatic arthritis, meta-analysis

Key messages:

- 1) The mean delay to diagnosis in axial SpA is 6.7 years worldwide.
- 2) This is significantly longer than the mean delay of 2.6 years reported in PsA studies.
- 3) Lower education, absence of extra-articular manifestations and younger onset age are associated with delay in axSpA.

Introduction

Axial spondyloarthritis (axSpA) is a chronic inflammatory disease characterised by significant inflammatory pain, stiffness and functional impairment [1]. Symptoms typically begin in early adulthood, which is a critical time for education, career, social networks and development of personal identity in general. Consequently, axSpA can significantly impact on mental health, quality of life and work productivity over the life course, at costs to the individual and the economy [2,3].

The disease impact is often compounded by a prolonged diagnostic delay, that is, time from onset of symptoms to getting a diagnosis. This may be explained by the lack of awareness of axSpA as a cause of back pain among patients and primary care providers and/or suboptimal referral to appropriate specialists or for investigations. Duration of delay is reported to range from 8 to 10 years – longer than many other rheumatic diseases - although estimates can vary considerably from study to study. Some studies have also found no improvement in diagnostic delay over recent decades [4], despite improved understanding of the disease and access to imaging.

There is abundant evidence that diagnostic delay is associated with worse functional impairment, greater radiographic progression, poorer quality of life and reduced response to treatment [5,6]. Those with longer delays to diagnosis also report greater work disability, unemployment and healthcare costs [5]. Although the impact of delay is well described, potential causes of delay (i.e., how delay can be improved) are not. Examining how delay durations vary across parts of the world and factors associated with delay will help inform targets for improvement.

The aims of this systematic review were to 1) describe global variation in diagnostic delay and 2) describe patient and disease factors that have been reportedly associated with delay to diagnosis. We also sought to 3) formally compare delay duration in axSpA with other SpA (e.g., psoriatic arthritis) to highlight the need and target for improvement.

Methods

We performed a systematic review and reported it in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [11]. The protocol for this review was pre-registered in advance (PROSPERO: CRD42020161887). We searched Medline, PubMed, EMBASE and Web of Science for relevant literature in September 2019 using the following search terms: (ankylosing OR spondyloarthritis OR psoriatic) AND ((delay AND diagnosis) OR (symptom AND (onset OR duration))). This search strategy was designed to include studies that describe diagnostic delay, even if delay was not their research objective. In response to peer review, bibliographies of all eligible studies and a prior systematic review [7] were also manually searched to identify additional titles.

Studies were included if they reported mean delay to diagnosis (i.e., the mean difference between age at symptom onset and diagnosis) or if they reported both mean age at onset *and* at diagnosis. Studies that defined eligibility using these variables were not eligible since estimates would not be representative. We excluded studies using the same (or very similar) cohort to studies already included. We also excluded studies reporting medians only. Letters and published conference abstracts were considered, as some prevalence studies may not be published as full articles but may have sufficiently detailed methodology and results. Non-systematic reviews, comments and editorials were excluded.

Two independent reviewers screened titles and abstracts, assessed full-texts for eligibility and extracted data from qualifying studies (BP, NH). Any discrepancy at each stage was resolved through discussion moderated by a third reviewer. Information from included studies was extracted into predefined tabulated summaries (Supplementary Table S1). Studies were assessed for risk of bias using adapted versions of the Newcastle Ottawa Scale (Supplementary Table S2).

Analysis

We pooled mean diagnostic delay using inverse variance weighted random-effects models (DerSimonian-Laird method). This was performed for studies of axSpA (including AS), then separately for psoriatic arthritis (PsA) and spondyloarthritis (SpA, which includes axSpA, PsA and other members of the SpA family). Where mean delay to diagnosis was not reported, it was imputed as the difference in mean age at symptom onset and mean age at diagnosis. Where the standard deviation of diagnostic delay was missing, we imputed it using methods recommended by Cochrane (in essence, based on standard deviations of age at onset, age at diagnosis and their correlation in all studies [8]) or the standard deviation of a study reporting the most similar mean delay duration. We

performed sensitivity analyses without imputed values. Further sensitivity analyses requested at peer review were performed by 1) restricting to studies using classification criteria only, and 2) excluding studies with potentially non-representative sampling (e.g., entirely male populations, or sample sizes of <30 that may give unstable population estimates). Heterogeneity of meta-analysis estimates was presented using the I^2 statistic. Funnel plots were used to assess risk of publication bias.

We used random-effects meta-regression to examine whether heterogeneity in axSpA diagnostic delay could be explained by study characteristics, i.e., year of publication (pre-2010, 2010-2015, post 2015), geography (regions defined by the World Health Organisation [9]), economic status of the country (World Bank economic class [10]), sample sources (e.g., single centre, multicentre etc), age at symptom onset (tertile) and proportion of males (tertile). Meta-regression was not performed for PsA and SpA due the limited number of studies. Analyses were performed using R version 3.6.2 and the “meta” and “metafor” packages.

Results

A total of 3286 publications were found from the literature search. 66 studies remained After excluding duplicates, irrelevant or ineligible studies, and studies using the same cohorts (or subsets thereof). A further 12 studies were found through manual bibliography searches. The study by Rojas-Vargas et al was excluded as it only included patients with ≤ 2 years of symptoms. The selection flowchart is shown in Supplementary Figure S1. The 78 included studies are summarised in Supplementary Table S1. 64 studies reported delays among axSpA patients, 8 PsA and 8 SpA. Feld et al [11] and Sørensen et al [12] reported delay in both axSpA and PsA. Bias scores were mostly 3 to 4 out of 6 stars (Supplementary Table S2 and Figure S2) indicating moderate bias.

Diagnostic delay in axSpA

Sample size for axSpA studies ranged from 5 to 2,887 patients. 47 studies were of AS (including 32 using modified New York criteria) and 17 of axSpA (including 11 using the ASAS criteria). Delay ranged from 2.8 years in a small Albanian study (of 54 cases over 6 years), to 11.1 years in a single UK centre [13,14]. The mean delay to diagnosis was 6.7 years overall (95% confidence interval 6.2 to 7.2, $I^2=99\%$) (**Figure 1**). The accompanying funnel plot is shown in Supplementary Figure S3.

Results of stratified meta-analysis are shown in **Table 1**. 39 axSpA studies were from countries in the European region, 9 West Pacific, 8 Eastern Mediterranean, 5 Americas and 3 South East Asia. Across these WHO regions, the mean delay and heterogeneity were not significantly different. When these studies were stratified according to World Bank economic class, the High-income group had longer mean delays than the upper- and lower-middle income countries. When mean delays were pooled according to country (with ≥ 3 studies), the average diagnostic delay was significantly shorter in Turkey and China than in the UK. Mean delay duration did not differ according to year of publication. Studies with older mean age of symptom onset showed trends for shorter delay durations.

WHO regions, economic class, recruiting centre, year of publication, age at symptom onset and male proportion were entered into a multivariable meta-regression model (Supplementary Table S3). Compared to high income countries, those in the upper- (by 2.5 years, $p<0.01$) and lower-middle income (3.7 years, $p=0.03$) category had shorter mean delay. Compared to studies from Europe, those from the Americas (by 2.1 years, $p=0.07$) and West Pacific region (2.9 years, $p=0.01$) had shorter mean delay.

Sensitivity analysis restricting to 43 studies using classification criteria showed similar mean delay duration of 6.5 years (Supplementary Figure S4). Diagnostic delay was 6.3 years (95%CI 5.6 to 7.0) for modified New York criteria AS and 7.1 years (95%CI 5.5 to 8.7) for ASAS criteria axSpA. Excluding

3 studies with potentially non-representative sampling (n<30 or all-male populations) did not change results (data not shown). Sensitivity analyses excluding studies with imputed mean and/or standard deviation of delay produced similar results (data not shown).




Figure 1. Pooled estimate of diagnostic delay in axial spondyloarthritis (including ankylosing spondylitis). Results ordered according to geography and year of publication. *Diagnostic delay

calculated from summary data for age at onset and diagnosis. **Standard deviation imputed for meta-analysis.

Table 1. Meta-analysis of delay duration stratified by study characteristics.				
	n	mean delay	95% CI	I ²
World Health Organisation regions				
European	39	7.09	6.46, 7.72	97.9%
West Pacific	9	5.92	4.34, 7.51	96.8%
Eastern Mediterranean	8	6.27	4.92, 7.61	97.7%
Americas	5	6.36	3.70, 9.03	99.8%
South East Asia	3	6.38	0, 14.2	96.7%
World Bank economic class				
High	39	7.56	7.04, 8.09	97.5%
Upper middle	20	5.37	4.54, 6.20	96.4%
Lower middle	5	5.59	3.21, 7.96	95.7%
Countries with ≥3 studies				
UK	9	8.65	7.35, 10.0	94.4%
Turkey	10	5.88	4.80, 6.96	90.3%
Italy	3	7.68	2.67, 12.69	99.6%
Iran	4	6.44	3.31, 9.58	98.2%
China	4	4.32	2.63, 6.00	79.1%
Recruiting methods				
Single centre	37	6.45	5.74, 7.16	99.0%
>1 centre	27	7.14	6.45, 7.84	99.2%
Year of publication				
<2010	12	6.75	5.66, 7.83	99.6%
2010-15	30	6.83	6.05, 7.61	97.1%
>2015	22	6.61	5.70, 7.53	99.3%
Age at symptom onset (tertiles)				
22.7 - 24.0 years	12	7.01	5.69, 8.33	97.7%
24.2 - 27.1 years	13	7.11	6.04, 8.19	98.5%
27.1 - 35 years	13	6.25	5.03, 8.23	99.0%
Proportion of males (tertiles)				
39-68%	20	6.95	6.08, 7.82	98.5%
69-80%	19	7.30	6.31, 8.29	98.2%
80-100%	19	5.65	4.91, 6.39	94.5%

Factors associated with delay to diagnosis

Most results were from unadjusted comparisons (**Table 2**). Delay was reportedly longer in males in studies by Bandinelli (10 v 6.3 years, $p=0.002$) and Sykes (9.4 v 8.3, $p=0.097$) [4,15], but longer in females in studies by Fallahi (8.7 v 7.7, $p=0.68$), Dincer (14 v 5.3, $p=0.06$), Hajjalilo (8.0 v 5.9, $p=0.14$), Jones (8.5 v 5.6) and Redeker (by 1.9 years, $p<0.05$) [16–20], albeit mostly not statistically significant. Similarly, 2 studies reported longer delay in those with peripheral arthritis [16,18], while 5 reported longer delays in those without [4,6,15,21,22]. There was also inconsistency in whether studies found HLA-B27 status to be associated with diagnostic delay: 4 studies reported significantly longer delays in HLA-B27 negative patients [16,17,20,23], while 5 other studies did not [15,21,22,24,25].

There was better consensus among the studies that longer delay was associated with: the absence of EAMs [4,18,24], lower education [16,17,21,26], and younger age of onset [20,21,25,26].

Table 2. Factors associated with longer delay to diagnosis in axial spondyloarthritis (results reported as mean duration in years).	
Aggarwal 2009 [24]	Absence EAMs v presence (8.7 vs 5.9, $p=0.03$) Onset <16 v >16 yrs (9.1 v 6.1, $p=0.03$)
Bandinelli 2016 [15]	Male v females (10 vs 6.3, $p=0.002$) Manual v non-manual workers (11 vs 8.3, $p=0.047$) Axial presentations compared to arthritis or enthesitis (9.0 vs 8.5 vs 4.3, $p=0.002$) Lower education (<high school v high school v university: 10 v 8.6 v 7.3, $p=0.076$)
Dincer 2008 [17]	HLA-B27 negative v positive (9.2 vs 5.3, $p=0.037$) Family history v none (10 vs 4.6 $p=0.003$) Onset \leq 16 v >16 yrs (8.9 v 5.5, $p=0.027$) Lower education (<9yrs v 9-11 v 12-13 v 14-15: 12 v 6.3 v 5.0 v 4.6, $p=0.018$) Females v males (14 v 5.3, $p=0.061$)
Fallahi 2016 [16]	Enthesitis v no enthesitis (8.8 vs 6.0, $p=0.007$) HLA-B27 negative v positive (10 vs 7.1, $p=0.013$) Lower education (correlation $r=0.24$ $p=0.002$) Presence of peripheral arthritis v absence (8.9 v 6.8, $p=0.086$)
Feldtkeller 2003 [23]	HLA-B27 negative v positive (11 v 8.5, $p<0.01$)
Gerdan 2012 [26]	With v without prior diagnosis of lumbar disc herniation (9.1 vs 6.2, $p=0.002$) First contact being rheumatology v non-rheumatology (8.1 vs 2.9, $p<0.001$) Younger age at onset ($b=-0.18$, $p=0.003$) Lower education ($b=-0.252$, $p=0.018$)
Hajjalilo 2014 [18]	Presence of peripheral arthritis v absence (11 vs 5.1 $p<0.001$) Absence of uveitis v presence (6.4 v 2.4 $p=0.02$) Presence of heel pain v absence (13 v 5.9 $p=0.004$) Females v males (8.0 v 5.9, $p=0.14$)
Jones 2014 [19]	Females v males (8.5 vs 5.6)

Masson Behar 2017 [21]	Univariable regression showed longer delay with Older age at diagnosis (b=0.15 p<0.001) Lower education (b=-1.7 p=0.03) Later calendar year of diagnosis (b=0.1 p=0.005) Multivariable regression showed longer delay with Older age at diagnosis (b=0.1, p<0.001) Entheseal pain v none (b=1.5 p=0.015) Absence of peripheral arthritis/dactylitis v presence (b=-1.7, p=0.005)
Nakashima 2016 [22]	Absence articular involvement vs presence (8.9 v 5.2, p=0.03) Disease onset pre-2000 v post (7.5 v 3.5 p=0.02)
Reed 2008 [27]	Delay longer with later calendar year and younger age at onset (p<0.05)
Seo 2015 [6]	Long-delay (v short delay <=8 years) category associated with: Absence of peripheral symptoms (OR 2.2, p=0.06) Prior diagnosis of mechanical back pain (OR 2.8, p=0.02) In univariate analysis, mechanical back pain remained significant in multivariable model
Sykes 2015 [4]	Absence of peripheral arthritis vs presence (9.4 v 7.6, p=0.045) Absence of IBD v presence (9.2 v 6.5, p=0.012) Presence of uveitis vs absence (10 v 8.4, p=0.033) Females v males (9.4 v 8.3, p=0.097)
Redeker 2018 (abstract) [20]	Multivariable regression showed longer delay in Female v males (b=1.9, 95%CI 1.1 2.7) Younger age of symptom onset, per 10yrs (-1.9, 95%CI -2.3, -1.5) HLA-B27 negative v positive (-3.6, 95%CI -5.1, -2.1) Psoriasis v no psoriasis (1.4, 95%CI 0.1, 2.7)
Resende 2018 (abstract) [25]	Presence of EAMs v absence (8.7 v 5.0, p<0.001) Younger age onset (r=-0.28, p<0.001)
EAM, extra-articular manifestations (anterior uveitis, psoriasis, inflammatory bowel disease)	

PsA and SpA

Sample size for 8 PsA studies ranged from 69 to 1970 patients. Diagnostic delay ranged from 1.0 years in the Dutch South-West Psoriatic Arthritis to 4.6 in a Swedish population-based cohort [28,29]. The mean delay to PsA diagnosis was 2.6 years (95%CI 1.6 to 3.6, I²=99%) (**Figure 2**).

Eight SpA studies ranged from 16 to 708 participants in size and 1.6 to 7.6 years in diagnostic delay. The mean delay to SpA diagnosis was 4.9 years (95%CI 3.3 to 6.6, I²=96%) (**Figure 2**). Funnel plot is shown in Supplementary Figure S3

Sensitivity analysis restricting to 3 PsA and 6 SpA studies using classification criteria showed similar results (Supplementary Figure S5).




Figure 2. Pooled estimate of diagnostic delay in psoriatic arthritis and spondyloarthritis. *Diagnostic delay calculated from summary data for age at onset and diagnosis. **Standard deviation imputed for meta-analysis.

Discussion

The mean delay to diagnosis was 6.7 years across 64 axSpA studies worldwide. Interestingly, countries classed as high-income by the World Bank had significantly longer delays to diagnosis than medium-income countries. Factors associated with delay to diagnosis varied and were often contradictory across studies; the most consistently reported factors were lower education, absence of extra-articular manifestations and younger age of onset. Diagnostic delay in axSpA was significantly longer than studies of PsA (2.6 years) and SpA (4.9 years).

Mean duration of delay varied significantly within (e.g., from 5.7 to 11 years in the UK) and between countries. This may reflect multiple factors that could not be assessed in this review, such as local healthcare infrastructure and awareness of the disease. Our finding that delay was longer in high-income countries was unexpected. It may be that research centres in these countries received referrals for the most diagnostically challenging cases or served comparably deprived areas. Conversely, it may be that only centres with good referral infrastructure are publishing research in middle-income countries.

Our meta-analysis showed no meaningful change in diagnostic delay over (publication) time. This is consistent with results from the UK [4,30], France [21] and Germany [20]. In stark contrast, delay to diagnosis improved dramatically in Japan (pre- v post-2000: 7.5 v 3.6 years [22]), Italy (1990s v 2000s: 7.4 v 2.1 years [31]), Egypt (pre- v post-2010: 11 v 4.6 years [32]) and Australia [27]. We could not examine the cause of this variation in detail, but diagnostic approaches likely varied from country to country. For example, the extent to which HLA-B27 and gender were associated with delay differed between countries, suggesting that these factors may have differential importance in their respective diagnostic process.

Inflammatory back pain in axSpA typically has an insidious onset, with subtle signs on clinical examination. There is also a plethora of highly prevalent differential diagnoses that may be incorrectly used to explain symptoms; for example, lumbar disc disease can co-exist with axSpA and prolong delay to diagnosis [6,26]. Peripheral joint involvement is relatively more acute in presentation, with clearer signs such as swelling and erythema. This may explain the much shorter diagnostic delay in PsA than in axSpA. Among axSpA studies, the presence of peripheral joint involvement was associated with shorter delay to diagnosis in Italian [15], UK [4], French [21] and Japanese [22] studies, while these patients had *longer* delays in Iran [16,18]. It may be the case that these Iranian patients were given other diagnoses prior to the correct axSpA label.

To reduce delay to diagnosis, a potential target would be to improve awareness of axSpA as a cause of back pain; general education was inversely associated with delay. Younger age of onset was also consistently associated with prolonged delay. (Although this may be an artefact of “delay” being derived from, and being dependent on, age at onset.) Education is needed among non-rheumatologists that axSpA is a cause of back pain in young people. However, there will be cases that remain more diagnostically challenging, such as patients with few SpA features.

A key strength of this review is the large and globally representative number of studies. Our unique search strategy allowed us to include studies that described diagnostic delay, even if delay was not their research objective. Such studies are less likely to be subject to subconscious bias from a prior delay hypothesis, thus their inclusion is a strength rather than weakness. There were however limitations. Diagnostic delay is known to be right-skewed in distribution, meaning that the mean is inflated above the median by a high proportion of people with disproportionately long delays. In other words, the mean may be sensitivity to these outliers (e.g., atypical clinical features or individuals with poor access to healthcare) and remain unchanged, even if diagnostic delay generally improved for many patients. We chose mean firstly because it permits meta-analysis, but also because median would take emphasis away from those with unusually long delays - precisely the individuals needing improvement to diagnosis. Some meta-analysis estimates for delay had negative lower-bounds in the confidence interval, which is not possible by definition. This is an artefact of the random-effects methodology; in each case, there is one study with a much shorter delay than others in the category, resulting in wide intervals required to cover the pooled estimate for this subgroup. This artefact disappears in fixed-effects models, which were not used in this study due to high heterogeneity between the studies. We did not review the impact of delay to diagnosis as this was recently reviewed [5]. Meta-regression examines relationships between summary data and should not be interpreted as traditional hypothesis testing of individual patient data. For example, proportion of males was not associated with diagnostic delay; this does not rule out a difference in delay between the sexes. Although most studies in our review did not report a statistically significant difference, a prior meta-analysis of SpA (excluding PsA) did [7]. Delay over time should also be interpreted with caution. Year of publication was the only available proxy for calendar time, since the recruitment period can be over many years and was often not reported. The intervals between recruitment and publication were generally homogenous in studies that did report this data.

Conclusion

The delay from symptom onset to diagnosis is 6.7 years on average in axSpA, which is significantly longer than 2.6 years for PsA. Although delay has improved over time in some parts of the world,

many countries such as the UK need additional efforts to improve delay to diagnosis. Lower education levels, absence of EAMs and younger age of onset were associated with longer delays; therefore, improved education for physicians and patients with back pain may help reduce diagnostic delay.

Declarations

Availability of data and materials: All data relevant to the study are included in the article or uploaded as online supplementary information.

Competing interests: none declared.

Funding: none.

Authors' contributions: SZ wrote the manuscript with significant contribution from all co-authors. BP, NLH, AEA performed the literature search, data extraction and quality assessment. SZ and DMH performed all statistical analysis and conceived of the project. All authors read and approved the final manuscript.

Acknowledgements: We thank Ms Karishma Nursiah for contributing to the design stage of the review, and the UK National Axial Spondyloarthritis Society (NASS) for their efforts to improve delay to diagnosis.

1. Sieper J, Braun J, Dougados M, Baeten D. Axial spondyloarthritis. *Nat Rev Dis Primer*. 2015 09;1:15013.
2. Martindale J, Shukla R, Goodacre J. The impact of ankylosing spondylitis/axial spondyloarthritis on work productivity. *Best Pract Res Clin Rheumatol*. 2015 Jun 1;29(3):512–23.
3. Zhao S, Thong D, Miller N, Duffield SJ, Hughes DM, Chadwick L, et al. The prevalence of depression in axial spondyloarthritis and its association with disease activity: a systematic review and meta-analysis. *Arthritis Res Ther*. 2018 Dec;20(1).
4. Sykes MP, Doll H, Sengupta R, Gaffney K. Delay to diagnosis in axial spondyloarthritis: are we improving in the UK? *Rheumatology*. 2015 Dec 1;54(12):2283–4.
5. Yi E, Ahuja A, Rajput T, George AT, Park Y. Clinical, Economic, and Humanistic Burden Associated With Delayed Diagnosis of Axial Spondyloarthritis: A Systematic Review. *Rheumatol Ther*. 2020 Mar;7(1):65–87.
6. Seo MR, Baek HL, Yoon HH, Ryu HJ, Choi H-J, Baek HJ, et al. Delayed diagnosis is linked to worse outcomes and unfavourable treatment responses in patients with axial spondyloarthritis. *Clin Rheumatol*. 2015 Aug;34(8):1397–405.
7. Jovaní V, Blasco-Blasco M, Ruiz-Cantero MT, Pascual E. Understanding How the Diagnostic Delay of Spondyloarthritis Differs Between Women and Men: A Systematic Review and Metaanalysis. *J Rheumatol*. 2017 Feb;44(2):174–83.
8. Higgins J, Li T, Deeks J. *Cochrane Handbook for Systematic Reviews of Interventions*. Chapter 6; Section 6.5.2.8: Imputing standard deviations for changes from baseline [Internet]. Version 6, 2019. [cited 2020 Feb 28]. Available from: <https://training.cochrane.org/handbook/current/chapter-06>
9. World Health Organisation. WHO regional offices. [Internet]. 2019. Available from: <http://www.who.int/about/regions/en/>
10. World Bank. List of economies 2017. [Internet]. 2019. Available from: databank.worldbank.org/data/download/site-content/CLASS.xls.
11. Feld J, Ye JY, Chandran V, Inman RD, Haroon N, Cook R, et al. Is axial psoriatic arthritis distinct from ankylosing spondylitis with and without concomitant psoriasis? *Rheumatology*. 2019 Oct 8;kez457.
12. Sørensen J, Hetland ML. Diagnostic delay in patients with rheumatoid arthritis, psoriatic arthritis and ankylosing spondylitis: results from the Danish nationwide DANBIO registry. *Ann Rheum Dis*. 2015 Mar;74(3):e12–e12.
13. Koko V, Ndrepepa A, Sknderaj S, Ploumis A, Backa T, Tafaj A. An Epidemiological Study on Ankylosing Spondylitis in Southern Albania. *Mater Socio Medica*. 2014;26(1):26.
14. Zhao SS, Radner H, Siebert S, Duffield SJ, Thong D, Hughes DM, et al. Comorbidity burden in axial spondyloarthritis: a cluster analysis. *Rheumatology*. 2019 Oct 1;58(10):1746–54.
15. Bandinelli F, Salvadorini G, Sedie AD, Riente L, Bombardieri S, Matucci-Cerinic M. Impact of gender, work, and clinical presentation on diagnostic delay in Italian patients with primary ankylosing spondylitis. *Clin Rheumatol*. 2016 Feb;35(2):473–8.

16. Fallahi S, Jamshidi AR. Diagnostic Delay in Ankylosing Spondylitis: Related Factors and Prognostic Outcomes. :7.
17. Dincer U, Cakar E, Kiralp MZ, Dursun H. Diagnosis delay in patients with ankylosing spondylitis: possible reasons and proposals for new diagnostic criteria. *Clin Rheumatol*. 2008 Apr;27(4):457–62.
18. Hajjalilo M, Ghorbanihaghjo A, Khabbazi A, Kolahi S, Rashtchizadeh N. Ankylosing Spondylitis in Iran; Late diagnosis and Its Causes. *Iran Red Crescent Med J [Internet]*. 2014 Apr 5 [cited 2020 Feb 22];16(4). Available from: <http://ircmj.com/en/articles/16025.html>
19. Jones A, Harrison N, Jones T, Rees JD, Bennett AN. Time to diagnosis of axial spondylarthritis in clinical practice: signs of improving awareness? *Rheumatology*. 2014 Nov 1;53(11):2126–7.
20. Redeker I, Callhoff J, Hoffmann F, Haibel H, Sieper J, Zink A, et al. COMORBID CONDITIONS ARE ASSOCIATED WITH HIGHER DISEASE ACTIVITY AND WORSE FUNCTIONAL STATUS IN AXIAL SPONDYLOARTHRITIS: A POPULATION-BASED ANALYSIS OF INSURANCE CLAIMS LINKED TO PATIENT SURVEY DATA. *Annals of the Rheumatic Diseases* 2019;78:1253-1254. In.
21. Masson Behar V, Dougados M, Etcheto A, Kreis S, Fabre S, Hudry C, et al. Diagnostic delay in axial spondyloarthritis: A cross-sectional study of 432 patients. *Joint Bone Spine*. 2017 Jul;84(4):467–71.
22. Nakashima Y, Ohishi M, Okazaki K, Fukushi J-I, Oyamada A, Hara D, et al. Delayed diagnosis of ankylosing spondylitis in a Japanese population. *Mod Rheumatol*. 2016 May 3;26(3):421–5.
23. Feldtkeller E, Khan M, van der Heijde D, van der Linden S, Braun J. Age at disease onset and diagnosis delay in HLA-B27 negative vs. positive patients with ankylosing spondylitis. *Rheumatol Int*. 2003 Mar;23(2):61–6.
24. Aggarwal R, Malaviya AN. Diagnosis delay in patients with ankylosing spondylitis: factors and outcomes—an Indian perspective. *Clin Rheumatol*. 2009 Mar;28(3):327–31.
25. Resende GG, Lage R, Malheiro O, Guimaraes D, Carvalho de Paula F, Carvalho D, et al. O20 DIAGNOSTIC DELAY IN SPONDYLOARTHRITIS: HOW CAN WE DO BETTER? XXXV Brazilian Congress of Rheumatology. *Adv Rheumatol* 58, 23 (2018).
26. Gerdan V, Akar S, Solmaz D, Pehlivan Y, Onat AM, Kisacik B, et al. Initial Diagnosis of Lumbar Disc Herniation Is Associated with a Delay in Diagnosis of Ankylosing Spondylitis. *J Rheumatol*. 2012 Oct;39(10):1996–9.
27. Reed MD, Dharmage S, Boers A, Martin BJ, Buchanan RR, Schachna L. Ankylosing spondylitis: an Australian experience. *Intern Med J*. 2008 May;38(5):321–7.
28. Vis M, Wervers K, Tchetverikov I, Kok MR, Korswagen L-A, van Groenendaal H, et al. Diagnostic Delay Leads to Worse Response to Treatment [abstract]. *Arthritis Rheumatol*. 2016; 68 (suppl 10). <https://acrabstracts.org/abstract/diagnostic-delay-leads-to-worse-response-to-treatment/>. Accessed February 28, 2020. :3.
29. Bremander A, Jacobsson LTH, Bergman S, Haglund E, Löfvendahl S, Petersson IF. Smoking is associated with a worse self-reported health status in patients with psoriatic arthritis: data from a Swedish population-based cohort. *Clin Rheumatol*. 2015 Mar;34(3):579–83.

30. Moran S, Longton C, Bukhari M, Ottewell L. AB0708 Delay To Diagnosis in Ankylosing Spondylitis: A Local Perspective *Annals of the Rheumatic Diseases* 2016;75:1146-1147.
31. Salvadorini G, Bandinelli F, Sedie AD, Riente L, Candelieri A, Generini S, et al. Ankylosing spondylitis: how diagnostic and therapeutic delay have changed over the last six decades. *Clin Exp Rheumatol*. 2012;30:561–5.
32. Abdelrahman F, Mortada M. AB0858 Impact of application of asas criteria for axial spondyloarthritis on the diagnostic delay in egyptian patients *Annals of the Rheumatic Diseases* 2018;77:1556-1557.