Assessing axSpA What You Need To Know

Understanding your axial spondyloarthritis (axSpA) and being involved in treatment decisions are key steps towards improved control of your disease.

Regular assessments are important to track and understand your disease activity. Knowledge about assessment tools can help you to have a more informed conversation with your doctor and make a difference in how you manage your axSpA.

How is axSpA disease activity measured?

Disease activity captures the level of some of the key symptoms experienced by patients, for example, pain, fatigue, joint swelling and morning stiffness as well as inflammation in your blood.¹

The most common measurements tools used by rheumatologists are the **Bath Ankylosing Spondylitis Disease Activity Index** (BASDAI) and the **Ankylosing Spondylitis Disease Activity Score** (ASDAS).²



BASDAI Bath Ankylosing Spondylitis Disease Activity Index

The BASDAI measures the activity of the disease based on your answers to six questions related to fatigue, spinal pain, peripheral joint pain/swelling, areas of localised tenderness, and duration and severity of morning stiffness.³



A higher BASDAI score reflects higher disease activity.⁴

If your BASDAI score is equal or above 4, it indicates that your disease is not adequately controlled and switching therapy could be discussed with your physician.⁴

ASDAS

Ankylosing Spondylitis Disease Activity Score



The ASDAS assessment includes three of the BASDAI questions (around spinal pain, duration of morning stiffness and joint pain/swelling), your overall assessment of your disease, and the objective level of inflammation, measured in your blood (C-reactive protein [CRP]).⁵

An ASDAS score of **under 2.1** is considered low disease activity and is an important target in disease control.^{6,7}

CRP

Between 2.1 and 3.5 is interpreted as high disease activity.^{6,7}

Over 3.5 is very high disease activity.^{6,7}



Inflammation can be measured through the

from your previous ASDAS score is considered a clinically important improvement, a change of at least 2.0 units is a "major improvement".⁶ These improvements are likely to be reflected in your overall quality of life.⁸ level of C-reactive protein (CRP) in your blood, which increases when there is inflammation in your body.⁹ A simple blood test can be done to check your CRP level.

Generally, CRP levels below 5 mg/L are considered normal.¹⁰

CRP is considered as part of the ASDAS measurement. Rheumatologists may also consider it alongside using the BASDAI.⁴



R

References: 1. Lukas C, Landewé R, Sieper J, Dougados M, Davis J, Braun J, van der Linden S, van der Heijde D; Assessment of Spondyloarthritis International Society. Development of an ASAS-endorsed disease activity socre (ASDAS) in patients with ankylosing spondylitis. Ann Rheum Dis. 2009 Jan;68(1): 18-24. doi: 10.1136/ard.2008.094870. Epub 2008 Jul 14. PMID: 18625618. 2. Klile G, Klile G, Kaike G, Nas K, Karkucak M, Capkin F, Dağin AZ, Çevik R, Özgörmen S. Comparison of ASDAS and BASDAI as a measure of disease activity in axial postratic arthritis. Clin Rheumatol. 2015 Mar;24(3):51-521. doi: 10.1007/s10067-014-27344. B. Epub 2014) Jul 2. PMID: 24985042. 3. Garrett S, Jenkinson T, Kennedy LG, Whitelock H, Gaisford P, Calin A. A new approach to defining disease status in ankylosing spondylitis: the Bath Ankylosing Spondylitis Disease Activity Index. J Rheumatol. 1994 Dec;21(12):2286-91. PMID: 7699630. 4. van der Heijde D, Ramiro S, Landewé R, Baraliakos X, Van den Bosch F, Sepriano A, Regel A, Ciurea A, Dagfinrud H, Dougados M, van Gaalen F, Géher P, van der Horst-Bruinsma I, Inman R, Jongkees M, Kilz U, Kwien T, Machado P, Marzo-Ortega H, Molto A, Navarro-Compàn V, Ozgocmen S, Pimentel-Santos F, Reveille J, Rudwaleit M, Sarejer J, Smapnio-Barros P, Wiek D, Braun J. 2016 update of the ASAS-ELULAR management recommendations for axial spondyloarthritis. Ann Rheum Dis. 2017 Jun;76(6):78-991. doi: 10.1136/ard.2016-210770. Epub 2017 Jan 13. PMID: 28087505. S. Sieper J, Rudwaleit M, Baraliakos X, Brandt J, Braun J, Burgos-Vargas R, Dougados M, Hermann K-G, Landewé R, Maksymowych W, van der Heijde D. The Assessment of SpondyloArthritis international Society C, ASAS Suppl 2:1114i44. doi: 10.1136/ard.2008.104018. PMID: 19433414. 6. Machado P, Landewé R, Lie E, Kvien TK, Braun J, Baker D, van der Heijde D. Assessment of SpondyloArthritis international Society. Ankylosing Spondyloits Disease Activity Socre (ASDAS): defining cut-off values for disease activity states and improvement scores. Ann Rheum Dis. 2011 Jan;70(1):47-



© UCB Biopharma SRL, 2023. All rights reserved. Date of preparation: April 2023 GL-N-DA-axSpA-2200116